



## Short Communication

Efficacy of Citrus Fruit Peel Extracts Against Pathogens Causing  
Gastrointestinal Disorders

NAREN DHIREN\*, CATHERINE LEO, SHARAN SARAVANAN AND MUTHU GANESH

Department of Biotechnology, Karpaga Vinayaga College of Engineering and Technology, Maduranthagam, TamilNadu-603308, India

## ARTICLE DETAILS

## Article history:

Received on 08 May 2013

Modified on 12 June 2013

Accepted on 20 June 2013

## Keywords:

Citrus peel,

Resistance and Prevention,

Gastrointestinal disorders.

## ABSTRACT

Gastrointestinal disorders are very common among adults; the Contaminated food and water are the main sources of the organisms which can cause gastric problems. Due to the intake of antibiotic drug against the pathogen causing gastrointestinal disorders, The bacterial strains are becoming resistant to the particular. To overcome this, citrus fruit peel extract can be used to treat these disorders. The citrus fruit peel extract used Citrus paradisi Macfad (Grapefruit) Citrus sinensis (Orange), Citrus limon (Lemon), and Citrus aurantifolia (Lime) gainst the gastrointestinal pathogens and lesser side effects than the synthetic drugs used against gastrointestinal pathogens.

© KESS All rights reserved

## INTRODUCTION

People have used citrus fruits as a source of medicines for thousands of years, but not in the citrus-flavored foods familiar to us today, like orange juice, lime pie or lemon slices on slabs of salmon. While we usually consume the flesh and nectar of these succulent fruits as food, herbalists have used the peels of these citrus fruit as medicine for numerous maladdies throughout history.

In some cases, these discoveries in the apothecary led to innovation in the kitchen. For example, in Asia, the use of orange zest, lemon zest and dried orange peel in cooking developed out of the knowledge of their application as remedies for digestive disorders. A little citrus peel in your diet can go a long way.

Food and waterborne disease is caused by eating contaminated food or drinking impure water. The gastric infections can be bacterial, viral, protozoal or helminthic. Many bacteria contaminating food and water can cause acute gastroenteritis or inflammation of the stomach and intestinal lining. Common symptoms of such food poisoning are nausea, vomiting and diarrhea. Traditional Chinese herbal medicine uses several citrus peels for specific health support, including those of mandarin orange (*Citrus reticulata* 'Blanco') and bitter orange (*C. aurantium*).

For hundreds of years, herbalists trained in Traditional Chinese Medicine (TCM) have used orange peel, known as chen pi or ju pi in Chinese medicine, to improve digestion, relieve intestinal gas and bloating, and resolve phlegm. This peel acts primarily on the digestive and respiratory systems. We apply it in conditions involving a sense of distension and fullness in the chest and upper middle abdomen combined with loss of appetite, vomiting or diarrhea, or coughs with copious phlegm.

Immature mandarin orange peel, known as qing pi in Chinese medicine, acts primarily on the liver and stomach to promote digestion, relieve food retention and abdominal distension, and promote good liver function. Practitioners of Chinese herbology use this herb when the sense of distension and discomfort lies primarily under the rib cage rather than the central abdomen.

Since the diverse bacterial introduction of antibiotics there has been tremendous increase in the resistance coexisting in the human body. The medicinal actions of citrus peels come in part from their primary essential oil, d-limonene. D-limonene has antimicrobial and anti-inflammatory properties. It also acts as a solvent for cholesterol, which has led some physicians to use it to dissolve cholesterol- D-limonene, neutralizes gastric acid and supports normal peristalsis, making it useful for relief of gastroesophageal reflux disease (GERD).

\*Author for Correspondence:

Email: narenbioresearch@gmail.com




Figure 1: Effect of ethanol extracts of the fruit peels on the organisms




Figure 2: Effect of crude extracts of the fruit peels on the organisms




Figure 3: Effect of acetone extracts of the fruit peels on the organisms




Figure 4: Effect of aqueous extracts of the fruit peels on the organisms




Figure 5: Effect of chloroform extracts of the fruit peels on the organisms




Figure 6: Effect of hexane extracts of the fruit peels on the organisms

## MATERIALS AND METHODS

The citrus fruit peels were procured from markets. They were washed well to remove dust and other external contaminants and by following standardized laboratory methods the Aqueous, Ethanol, Acetone, Chloroform, Hexane and Crude extracts were prepared. The Antibacterial Activity of the Gram Negative Microorganisms selected for this study (*E. coli*, *Salmonella* spp and *Shigella* spp) was conducted using Agar Well Method (Cup Plate Method) and Filter Paper Method. Kirby Bauer Method was employed for Antibiotic Sensitivity Testing. intestinal disorders than the antibiotics usually administered.

### Activity Assessment:

The effect of four citrus fruit peels (using aqueous, crude and solvents like ethanol, acetone, chloroform and hexane) on three common pathogens which causes stomach disorders were studied. In addition, a comparative study with antibiotics and various plant extracts was also carried out. Following the incubation of fruit peel extracts with the organisms, the zone of inhibition was measured in mm. The following results as per the inhibition were thus obtained. In the ethanol extract concentrate, *Citrus aurantifolia* and *Citrus limon* showed a better inhibitory effect on *E. coli* and *Shigella* spp. *Salmonella* showed less inhibition compared to other two and *Citrus paradisi* Macfad, showed less inhibition to *Shigella* and *Salmonella* spp. *Citrus sinensis* showed inhibitory effect on *E. coli* and *Shigella* spp, than *Salmonella*. In crude extract concentrate, *C. aurantifolia* showed more inhibitory activity against *Shigella* spp than *E. coli* and *Salmonella*. In *C. limon* inhibitory activity was seen more in *E. coli* compared to *Shigella* spp, and *Salmonella*. In *C. sinensis*, less zone of inhibition was witnessed than in others. In acetone extract concentrate, *C. aurantifolia* showed better inhibition against *E. coli* and *Shigella* spp. In *C. paradisi* Macfad less inhibition was seen in all three organisms. In aqueous extract, *C. sinensis* inhibitory activity was better in *E. coli* but less in *Salmonella* spp. *C. aurantifolia* showed inhibitorier activity against *Shigella* spp and *E. coli* than *Salmonella* spp. *C. paradisi* Macfad showed inhibitory activity against all the organisms. None of the chloroform extracts showed any result for any of the organisms. For hexane extract concentrate, *C. sinensis* showed inhibitory effect in *E. coli* while for other two organisms it didn't show much

effect. *C. aurantifolia* and *C. limon* showed inhibitory activity against *E. coli* than the other organisms. *C. paradisi* Macfad did not show much significant effect on the organisms (Figure 1-6).

## CONCLUSION

Microorganisms which cause infections are becoming resistant to the drugs that are being commercially used against pathogens causing gastrointestinal disorders. This citrus fruit peel extract study against the common pathogens of the gastrointestinal tract revealed them as better inhibitory agents than synthetic compounds. There is the need for novel compounds which can have a significant effect on these organisms and it is always preferred to take natural medicines which have lesser side effects than the synthetic ones.

## REFERENCES

- [1] Flavor and Extract Manufacturers' Association D-Limonene Monograph, 1-4, Flavor and Extract Manufacturers' Association; Washington, DC: 1991.
- [2] Chutia, M., Deka Bhuyan, P., Pathak, M.G. and Sarma T.C. 2009. Antifungal activity and chemical composition of *Citrus reticulata* Blanco essential oil against phytopathogens from North East India. *Food Sci. Technol.* 42(3):777-780.
- [3] Nannapaneni, R. Muthaiyan, A., Candall, P.G., Johnson, M.G., O'Bryan, C.A., et al., 2008. Antimicrobial activity of commercial Citrus-based natural extracts against *Escherichia coli* O15 isolates and mutant strains. *Foodborn. Pathog. Dis.* 5(5):695-699.
- [4] Schloegel, M. and Dazak, P., 2004. Conservation medicine: tackling the root cause of emerging infectious diseases and seeking practical solutions. *Wildlife Track.* 8(4):1-7.
- [5] Tomatake, H., Koga, T., Yamato, M., Kassu, A. and Ota, F. 2006. Antibacterial activity of citrus fruit juices against *Vibrio* species. *J. Nutr. Sci. Vitaminol.* 52(2):17-160.
- [6] The United States Code of the Federal Regulations, Title 21, Part 182.60.
- [7] Hakim IA, Harris RB, Ritenbaugh C. Citrus peel use is associated with reduced risk of squamous cell carcinoma of the skin. *Nutr Cancer* 2000; 37:161-168.
- [8] Igimi H, Nishimura M, Kodama R, Ide H. Studies on the metabolism of d-limonene

- (p-mentha-1,8-diene). I. The absorption, distribution and excretion of d-limonene in rats. *Xenobiotica* 1974; 4:77-84.
- [9] Kodama R, Yano T, Furukawa K, et al. Studies on the metabolism of d-limonene (p-mentha-1, 8-diene). IV. Isolation and characterization of new metabolites and species differences in metabolism. *Xenobiotica* 1976; 6:377-389.
- [10] Crowell PL, Lin S, Vedejs E, Gould MN. Identification of metabolites of the antitumor agent d-limonene capable of inhibiting protein isoprenylation and cell growth. *Cancer Chemother Pharmacol* 1992; 31:205-212.